

### GUARANTEED CONTENT

	W/W
Total Nitrogen (N)	8%
Ammonium Nitrogen (NH <sub>4</sub> -N)	3%
Urea Nitrogen (NH <sub>2</sub> -N)	5%
Soluble in Water and Neutral Ammonium Citrate	
Phosphorus Pentaoxide (P <sub>2</sub> O <sub>5</sub> )	11%
Water Soluble Phosphorus Pentaoxide (P <sub>2</sub> O <sub>5</sub> )	10%
Water Soluble Potassium Oxide (K <sub>2</sub> O)	15%
Water Soluble Magnesium Oxide (MgO)	2 %
Water Soluble Boron (B)	0,3%
Water Soluble Zinc (Zn)	0,04%

- It meets the magnesium needs of plants thanks to the 2% magnesium it contains.
- Nitrogen is in a form that is DCDA chelated, does not wash out and can remain in the soil for 70 - 110 days.
- Its phosphorus can be absorbed quickly and is completely soluble in soil.
- Potassium does not have any problems with washing or dissolving.
- Since it is in completely organic form, it increases the amount of organic matter in the soil



GARANTİ EDİLEN İÇERİK	% (w/w)
Totale Azot (N)	% 8
Ammoniyum Azot (NH <sub>4</sub> -N)	% 3
Üre Azotu (NH <sub>2</sub> -N)	% 5
Neutrale Ammoniyum Sitrat ve Suda Çözünür Fosfor Pentoksit (P <sub>2</sub> O <sub>5</sub> )	% 11
Suda Çözünür Fosfor Pentoksit (P <sub>2</sub> O <sub>5</sub> )	% 10
Suda Çözünür Potasyum Oksit (K <sub>2</sub> O)	% 15
Suda Çözünür Magnezyum Oksit (MgO)	% 2
Suda Çözünür Bor (B)	% 0,3
Suda Çözünür Çinko (Zn)	% 0,04

### KULLANIM YERİ, ZAMANI VE DOZU

CROP	APPLICATION	SOI APPLICATION
<b>ALL GREENHOUSE AND GREENHOUSE VEGETABLES</b>	3 - 4 applications from planting to harvest	2 - 3 kg/da
<b>ALL OPEN FIELD VEGETABLES</b>	3 - 4 applications from planting to harvest	2 - 3 kg/da
<b>MELON, WATERMELON AND PUMPKİN</b>	2 - 3 applications from planting to the beginning of ripening	2 - 3 kg/da
<b>POTATO, ONİON, RADİSH, CARROT, GARLIC</b>	2 - 3 applications from planting to the beginning of ripening	1 - 1,5 kg/da
<b>STRAWBERRY</b>	3 - 4 applications from vegetative period to harvest	2 - 3 kg/da
<b>TUBEROUS PLANTS (POTATO, CARROT, SUGAR BEET, ONİON ETC.)</b>	It is applied throughout the season from germination.	1,5-2 L / da
<b>BANANA</b>	Starting from 2-3 applications 30-45 days before birth (April-May).	2 - 3 kg/da
<b>VİNEYARD</b>	3 - 4 applications before flowering until harvest	1 - 1,5 kg/da
<b>APPLE, PEAR, CHERRY, SOUR CHERRY, APRİCOT, ALMOND, HAZELNUT, WALNUT, PEACH, PİSTACHİO ETC.</b>	Until the petals stand upright	2,5 - 3 kg/da
<b>FİG, OLİVE AND CİTRUS</b>	3 - 4 applications before flowering until harvest	2,5 - 3 kg/da
<b>ALL INDUSTRIAL AND FIELD CROPS</b>	3 - 4 applications before flowering until harvest	2,5 - 3 kg/d