


## **NPK FORMULATIONS**

20-20-20 + TE
18-18-18 + TE
12-6-32 + TE
12-32-6 + TE



- Formulated with low EC and Low pH rate raw material
- · Can be used safely for various crops.

26

- · Can be apply directly to the soil, by drip irrigation
- Average Application Rate is during growth stage 5-30 kg/ha/day dose can be applied. Application rates may differ according to plant variety and necessity.

## **RECOMMENDATION FOR APPLICATION:**

CROP	SOİL APPLİCATİON	FOLİAR	APPLICATION TIME
GREENHOUSE AND OPEN FİELD VEGETABLES	1 – 3kg/Da	300 – 500 gr	Applied at 15 to 20-day intervals after initial fruits
WİNTER VEGETABLES WİTH EDİBLE LEAVES	1 – 3kg/Da	350 – 500 gr	Applied at 15 to 20-day intervals starting from when plant reaches 3 or 4-leaf period.
TUBER PLANTS	1 – 3kg/Da	250 – 300 gr	Applied at 15 to 20-day intervals starting from when plant reaches 4 or 6-leaf period and after initial fruits.
ALL FRUİT TREES	1 – 4kg/Da	400 – 600 gr	Applied 2 to 3 times with the onset of growth.
INDUSTRIAL PLANTS	1 – 3kg/Da	300 – 400 gr	Applied at 15 to 20day intervals starting from when plant reaches 5 or 6-leaf period.
ALL LEGUMES	1 – 3kg/Da	300 – 400 gr	Applied at 15 to 20day intervals starting from when plant reaches 5 or 6-leaf period.
GRAİNS	-	250 – 300 gr	Applied during tillering period
GREEN AREAS	-	250 – 300 gr	Applied every two weeks starting from early growth period.

Fertiline

20-20-20

Fertiline

12-6-32

Fertiline

18-18-18