



GUARANTEED CONTENT

Organic Matter
Humic Acid + Fulvic Acid
Water Soluble Potassium Oxide (K₂O)
pH

% (w/w)

: % 8
: % 15
: % 1,5
: 10-12

- By regulating soil conditions, it increases root development.
- Detox helps to benefit from the nutrients that are connected to the soil.
- Prevents wash off in fertilizer applications.
- Provides increased structure and micro-organism activity of the soil.
- Prevents salinity in the soil.



| CROP | APPLICATION TIME | SOIL APPLICATION | FOLIAR |
|---|---|------------------|--------------|
| TOMATO, PEPPER, EGGPLANT | Applied at 20-day intervals starting from 10 to 15 days after planting until the end of harvest | 500 – 700 cc | 200 cc |
| CUCUMBER, MELON, SQUASH WATERMELON, | Applied during branching, fruit setting period and at 20 day-intervals afterwards | 500 – 700 cc | 200 – 250 cc |
| POTATO, ONION, LETTUCE, SPINACH, CARROT, PARSLEY, CABBAGE, | At 20-day intervals after plants reach 2- or 3-leaf period | 500 – 700 cc | 200 – 300 cc |
| BEAN, CHICKPEA, BROAD BEAN | Applied at 15-day intervals after plants reach the height of 5 to 10 cm until inflorescence period, and then at 20-day intervals until the end of harvest | 500 cc | 200 – 300 cc |
| CITRUS FRUITS | Applied at 20-day intervals starting from new bud and leaf growth. | 500 – 800 cc | 200 – 300 cc |
| APPLE, PEAR, QUINCE, APRICOT, PEACH, PLUM, CHERRY, ALMOND, OLIVE | Applied at 20 to 30 day-intervals before inflorescence and after fruit setting periods. | 100 cc/tree | 500 – 800 cc |
| STRAWBERRY | Applied at 15 to 20-day intervals starting from early vegetation. | 500 – 700 cc | 200 – 300 cc |
| VINEYARD | Applied when offshoots reach 10 to 15 cm, before inflorescence, and throughout seed setting and growth periods. | 700 – 900 cc | 200 – 300 cc |
| WHEAT, BARLEY, PADDY | Applied during tillering, bolting and earing periods. | 500 – 700 cc | 250 – 350 cc |
| CORN | Applied before planting or when plants reach the height of 10 to 15 cm and 8- to 10-leaf periods until the start of harvest period. | 800 – 1000 cc | 250 – 350 cc |
| COTTON | Applied at 20 to 25 day-intervals when plants reach 3- to 5-leaf period during bolling and later periods | 500 – 700 cc | 250 – 400 cc |
| ORNAMENTAL PLANT | Applied at 15 to 20-day intervals throughout vegetation period. | 800 – 1000 cc | 250 – 350 cc |