



### GUARANTEED CONTENT

Total Organic Matter  
Alginic Acid  
Water Soluble Potassium Oxide (K<sub>2</sub>O)  
EC  
pH

### % (w/w)

: % 5  
: % 0,2  
: % 5  
: 10.3 dS/m  
: 8-10

- Due to alginic acid content in the formulation, it is efficient on soil and plant.
- When foliar applied, it increases uptake of plant nutrients.
- Bella regulates soil pH. It increases water holding capacity.
- Supports root development and increases capillary root.
- Fastens uptaken of plant nutrients from roots to plants.
- Contains in different levels of plant growth compounds, mineral products and carbohydrates.

### RECOMMENDATION FOR APPLICATION:

| CROP                    | APPLICATION TIME  | FOLIAR DOSE    |
|-------------------------|---|----------------|
| <b>COTTON</b>           | the first application at 4-6-leaf stage. Follow up with applications, 2-3 times fortnightly   | 200-250 cc/da  |
| <b>GRAPES</b>           | 3 to 4 times during the growing season.   | 150 -200 cc/da |
| <b>CİTRUS</b>           | 2-3 application from beginning flowering  | 250-300 cc/da  |
| <b>SUGARBEET</b>        | 2 application during the growing season.  | 150 -200 cc/da |
| <b>VEGETABLES</b>       | 2-3 application from beginning flowering, fortnightly   | 150 -200 cc/da |
| <b>FRUIT TREES</b>      | 2-3 application from starting shoot development, fortnightly  | 200-250 cc/da  |
| <b>POTATOES</b>         | Apply 14 day intervals during periods of vegetative growth  | 150 -200 cc/da |
| <b>CEREALS</b>          | 2 application tillering and stem elongation   | 150 -200 cc/da |
| <b>GREENHOUSE</b>       | 2-4 application from planting to harvest  | 200-250 cc/da  |
| <b>MAİZE</b>            | 2 application during the growing season   | 200-250 cc/da  |
| <b>STRAWBERRY</b>       | 3-4 application intervals 20-30 days from planting  | 150 -200 cc/da |
| <b>GREEN FIELD</b>      | from spring at intervals of 20-30 days  | 150 -200 cc/da |
| <b>SEED APPLICATION</b> | Depending on the seed size, 5 cc of BELLA is added to 1 liter of water and the seeds are kept for 1 hour. Planting is done after it dries slightly. |                |

